



NT@UW-06-08
JLAB-THY-06-484
UNH-06-02

The Gell-Mann – Okubo Mass Relation among Baryons from Fully-Dynamical Mixed-Action Lattice QCD

Silas R. Beane,^{1,2} Kostas Orginos,^{2,3} and Martin J. Savage⁴

(NPLQCD Collaboration)

¹*Department of Physics, University of New Hampshire, Durham, NH 03824-3568.*

²*Jefferson Laboratory, 12000 Jefferson Avenue, Newport News, VA 23606.*

³*Department of Physics, College of William and Mary, Williamsburg, VA 23187-8795.*

⁴*Department of Physics, University of Washington, Seattle, WA 98195-1560.*

Abstract

We explore the Gell-Mann–Okubo mass relation among the octet baryons using fully-dynamical, mixed-action (domain-wall on rooted-staggered) lattice QCD calculations at a lattice spacing of $b \sim 0.125$ fm and pion masses of $m_\pi \sim 290$ MeV, 350 MeV, 490 MeV and 590 MeV. Deviations from the Gell-Mann–Okubo mass relation are found to be small at each quark mass.

I. INTRODUCTION

A decade before the formulation of Quantum Chromo-Dynamics (QCD) as the theory of the strong interactions, Gell-Mann [1] and Ne’eman [2] proposed that the low-lying hadrons could be embedded into irreducible representations of the flavor group $SU(3)$, with a fundamental representation containing three quarks: up, down and strange. This provided a relatively simple theory for understanding the plethora of strongly-interacting particles that had been seen at that time, although it was a radical proposal as the quarks had not been seen in isolation. Now that QCD is known to be the theory of the strong interaction, classification of the strongly-interacting particles by their transformation properties under the flavor group $SU(3)$ (and the isospin subgroup $SU(2)$) follows naturally. Of course, the flavor symmetries are only approximate, as they are broken by the differences in the quark masses. The impact that these mass differences have on hadronic spectra and interactions is determined by how they compare to the scale of strong interactions. The $SU(2)$ symmetry of isospin is a very-good symmetry of nature, being violated only at the percent level by both the electromagnetic interactions and the mass difference between the up and down quarks. The approximate $SU(3)$ symmetry is violated typically at the $\sim 30\%$ level due to the large mass difference between the strange quark and the up and down quarks as compared to the scale of the strong interaction. However, there are instances where $SU(3)$ symmetry works much better than this, as in the case of matrix elements of an $SU(3)$ charge operator between states in the same irreducible representation, which are protected by the Ademollo-Gatto theorem [3].

In addition to providing a well-defined scheme for understanding the spectrum of hadrons, flavor symmetry also provides a vital tool for the classification of operators, such as those responsible for nonleptonic decays. It is well established that the octet component of the nonleptonic weak interactions dominates over the **27** component, a feature known as octet dominance. This is a generic feature of strong interactions: the contributions from higher-dimensional representations of $SU(3)$ are suppressed compared to those from lower-dimensional representations. The origin of such suppressions is not well understood, but is found in perturbative calculations, and also by consideration of the large- N_C limit of QCD. The Gell-Mann–Okubo mass relation among the baryons is a classic example of this feature of QCD. When all three quark masses are equal, the light-quark mass matrix transforms as a singlet under flavor $SU(3)$, and as the strong interactions are also flavor singlets, higher-dimensional representations of $SU(3)$ are not induced, and

$$M_N = M_\Sigma = M_\Lambda = M_\Xi \quad , \quad (1)$$

where $N = p, n$ denotes the nucleon isodoublet, $\Sigma = \Sigma^+, \Sigma^0, \Sigma^-$ denotes the Σ isotriplet, and $\Xi = \Xi^-, \Xi^0$ denotes the Ξ isodoublet. In the limit of exact isospin symmetry, but broken $SU(3)$ symmetry, the light-quark mass matrix takes the form

$$m_q = \begin{pmatrix} \overline{m} & 0 & 0 \\ 0 & \overline{m} & 0 \\ 0 & 0 & m_s \end{pmatrix} \quad , \quad (2)$$

where $\overline{m}$ denotes the mass of the up and down quarks, while m_s denotes the mass of the strange quark. The mass matrix m_q transforms as an $\mathbf{8} \oplus \mathbf{1}$ under $SU(3)$ transformations. A single insertion of m_q in an octet baryon mass, or, more generally, an insertion in the

bm_l	m_π (MeV)	m_π/f_π	$10^3 \times T$ (l.u.)	octet centroid (l.u.)	$10^3 \times \delta_{\text{GMO}}$
0.007	293.1 ± 1.5	1.983 ± 0.020	2.4 ± 1.8	0.8128 ± 0.0082	3.0 ± 2.3
0.010	354.0 ± 0.9	2.332 ± 0.012	3.3 ± 1.0	0.8218 ± 0.0080	4.1 ± 1.3
0.020	492.6 ± 1.1	3.077 ± 0.014	0.34 ± 0.56	0.8724 ± 0.0055	0.39 ± 0.64
0.030	591.8 ± 0.9	3.588 ± 0.011	0.07 ± 0.42	0.9162 ± 0.0041	0.07 ± 0.46

TABLE I: Violations of the Gell-Mann–Okubo mass relation computed with lattice QCD. The first and second columns list the sea-quark masses employed in this work and the corresponding pion masses in MeV, respectively. We have used $b = 0.125$ fm to set the scale. The fourth column corresponds to T defined in eq. (3), measured in lattice units (l.u.), which is also the numerator of eq. (7). The fifth column corresponds to the location of the centroid of the lowest-lying baryon octet (also in lattice units), which is the denominator of eq. (7). All errors quoted are statistical. The systematic errors due to fitting are all significantly smaller than the statistical errors and thus are not shown.

baryon mass that transforms as an **8** under $SU(3)$ ¹, splits the masses of the baryons, but leaves one relation among the masses intact,

$$T \equiv M_\Lambda + \frac{1}{3}M_\Sigma - \frac{2}{3}M_N - \frac{2}{3}M_\Xi = 0 \quad , \quad (3)$$

the Gell-Mann-Okubo (GMO) mass relation [1, 4]. Multiple insertions of m_q , either through operators in the low-energy effective field theory, or through loops involving the light mesons, or both, will lead to violations of this relation. As T vanishes for $SU(3)$ breaking that transforms as $\mathbf{8} \oplus \mathbf{1}$, a non-zero value of T can result only from contributions transforming as **27**.

The typical scale of $SU(3)$ breaking is set by the differences in the squares of the meson masses,

$$Q \sim \frac{m_K^2 - m_\pi^2}{\Lambda_\chi^2} \sim 0.23 \quad , \quad (4)$$

where $\Lambda_\chi \sim 1$ GeV. The mass differences between baryons in the octet that differ in strangeness by one unit are then expected to scale as $\Delta M \sim Q\Lambda_\chi$, and one naively expects violations of the GMO relation to be $T \sim Q^2\Lambda_\chi \sim 52$ MeV. Experimentally, at $(m_\pi/f_\pi)^{\text{expt}} = 1.045$, one finds that $T^{\text{expt}} = 8.76 \pm 0.08$ MeV², where the isomultiplet averages of the masses have been used in eq. (3). Some theoretical understanding of why the GMO mass relation works better than naive expectations would suggest can be gained by considering the large- N_C limit of QCD, where it is found that violations to this relation are further suppressed by a factor of $1/N_C$ [5] giving $T \sim Q^2\Lambda_\chi/N_C \sim 17$ MeV, which is close to the experimental value.

Given the smallness of the violation of the GMO relation found experimentally, it is natural to ask whether this smallness persists at other values of the light-quark masses,

¹ Such a contribution will arise from n insertions of m_q , i.e. resulting from the **8**'s in the product $[\mathbf{1} \oplus \mathbf{8}] \otimes [\mathbf{1} \oplus \mathbf{8}] \otimes \dots \otimes [\mathbf{1} \oplus \mathbf{8}]$.

² The electromagnetic contribution to the violation of the GMO mass relation is expected to be an order of magnitude less than T^{expt} .

or is no more than an accident for the physical values of the quark masses. In this work we address this issue by computing deviations from the GMO mass relation at a number of different light-quark masses. Following the LHP collaboration (LHPC) [6, 7], our computation is a hybrid lattice QCD calculation using domain-wall valence quarks from a smeared-source on three sets of $N_f = 2 + 1$ asqtad-improved [8, 9] MILC configurations generated with rooted-staggered sea quarks [10]. In the generation of the MILC configurations, the strange-quark mass was fixed near its physical value, $bm_s = 0.050$, (where b is the lattice spacing) determined by the mass of hadrons containing strange quarks. The two light quarks in the four sets of configurations are degenerate (isospin-symmetric), with masses $bm_l = 0.007, 0.010, 0.020$ and 0.030 . Some of the domain-wall valence propagators were previously generated by LHPC on each of these sets of lattices. The domain-wall height is $M_5 = 1.7$ and the extent of the extra dimension is $L_5 = 16$. The parameters used to generate the light-quark propagators have been “tuned” so that the mass of the pion computed with the domain-wall propagators is equal (to few-percent precision) to that of the lightest staggered pion computed with the same parameters as the gauge configurations [10]. The MILC lattices were HYP-blocked [11] and Dirichlet boundary conditions were used to reduce the time extent of the MILC lattices from 64 to 32 time-slices in order to save time in propagator generation. Various parts of the lattice were employed to generate multiple sets of propagators on each lattice. We analyzed three sets of correlation functions on 564 lattices with $bm_l = 0.030$, 486 lattices with $bm_l = 0.020$, 658 lattices with $bm_l = 0.010$, and 468 lattices with $bm_l = 0.007$. The lattice calculations were performed with the *Chroma* software suite [12, 13] on the high-performance computing systems at the Jefferson Laboratory (JLab).

We denote as $C_X(t)$ the correlator as a function of time slice for baryon X , with large-time behavior

$$C_X(t) \rightarrow e^{-M_X t} . \quad (5)$$

We found that the deviations from the GMO mass relation were determined most precisely by forming the products and ratios of correlation functions associated with each of the baryons in the octet,

$$G^{\text{GMO}}(t) = \frac{C_\Lambda(t)}{C_N(t)^{2/3}} \frac{C_\Sigma(t)^{1/3}}{C_\Xi(t)^{2/3}} \rightarrow e^{-(M_\Lambda + M_\Sigma/3 - 2M_N/3 - 2M_\Xi/3)t} . \quad (6)$$

The resulting correlation functions are shown in fig. 1 and corresponding effective mass plots are shown in fig. 2. It is convenient to define a normalization factor with which to measure deviations from the GMO mass relation. We normalize the deviation to the centroid of the baryon octet, and define the fractional violation of the GMO mass relation, δ_{GMO} , to be

$$\delta_{\text{GMO}} = \frac{M_\Lambda + \frac{1}{3}M_\Sigma - \frac{2}{3}M_N - \frac{2}{3}M_\Xi}{\frac{1}{8}M_\Lambda + \frac{3}{8}M_\Sigma + \frac{1}{4}M_N + \frac{1}{4}M_\Xi} . \quad (7)$$

The experimental deviation translates into: $\delta_{\text{GMO}}^{\text{expt}} = 0.00761 \pm 0.00007$. The results of our fully-dynamical lattice QCD calculation are shown in Table I, and are plotted in fig. 3. The computation shows that the deviations from the GMO mass relation are small, and in fact less than the experimental value, at all quark masses accessible to us. It is now quite clear that the small violation of the GMO relation observed experimentally is not an accident




FIG. 1: *The logarithm of the correlation function $G^{\text{GMO}}(t)$ for the four quark masses employed in this work. Each has been off-set vertically for display purposes.*

due to the physical values of the light-quark masses, but is a generic feature of the baryon spectrum.

Heavy-baryon chiral perturbation theory [14, 15] (HB χ PT) can be used to compute corrections to the GMO mass relation. The leading-order correction arises at one-loop in the chiral expansion, and is dictated by the mass differences between the pseudo-Goldstone bosons, the π , K and η . It is found that [16, 17, 18]

$$M_\Lambda + \frac{1}{3}M_\Sigma - \frac{2}{3}M_N - \frac{2}{3}M_\Xi = \frac{1}{24\pi f_\pi^2} \left[\left(\frac{2}{3}\mathcal{D}^2 - 2\mathcal{F}^2 \right) (4m_K^3 - 3m_\eta^3 - m_\pi^3) - \frac{\mathcal{C}^2}{9\pi} (4F_K - 3F_\eta - F_\pi) \right], \quad (8)$$

where the function $F_c = F(m_c, \Delta, \mu)$ is

$$F(m, \Delta, \mu) = (m^2 - \Delta^2) \left(\sqrt{\Delta^2 - m^2} \log \left(\frac{\Delta - \sqrt{\Delta^2 - m^2 + i\epsilon}}{\Delta + \sqrt{\Delta^2 - m^2 + i\epsilon}} \right) - \Delta \log \left(\frac{m^2}{\mu^2} \right) \right) - \frac{1}{2}\Delta m^2 \log \left(\frac{m^2}{\mu^2} \right), \quad (9)$$

and we hold the Delta-Nucleon mass-splitting, Δ , fixed at its physical value, $\Delta^{\text{expt}} =$




FIG. 2: Effective mass plots for the correlation function $G^{\text{GMO}}(t)$ at the four quark masses employed in this work.




FIG. 3: Deviations from the Gell-Mann–Okubo mass relation for the lowest-lying baryon octet. The red diamond corresponds to the experimental value. The brown band corresponds to the leading-order one-loop $SU(3)$ HB χ PT prediction, and its expected $\sim 30\%$ uncertainty. The black data points are the results of our fully-dynamical lattice QCD calculation at pion masses of $m_\pi \sim 293$ MeV, 354 MeV, 493 MeV and 592 MeV.

293 MeV³. To the order we are working, the μ independence of the baryon GMO relation, eq. (8), is protected by the meson GMO relation, and thus no counterterms are required. For vanishing decuplet-octet mass splitting, the function F tends to $F_P \rightarrow \pi m_P^3$. The constants $\mathcal{D}$ and $\mathcal{F}$ are the axial couplings between the octet baryons and the octet of pseudo-Goldstone bosons, while the constant $\mathcal{C}$ is the axial coupling among the decuplet of baryon resonances, the octet of baryons and the octet of pseudo-Goldstone bosons. At tree-level these couplings take the values [19] $\mathcal{F} = 0.5$, $\mathcal{D} = 0.8$ and $\mathcal{C} = 1.5$, while at one-loop level they take the values [19] $\mathcal{F} = 0.4$, $\mathcal{D} = 0.6$ and $\mathcal{C} = 1.2$. In fig. 3 we have shown (the brown band) the one-loop prediction of HB χ PT, which is seen to agree, within uncertainties, with both the experimental value, and also the results of our lattice calculation. In obtaining this curve, we have divided the HB χ PT expression of eq. (8) by the experimental value of the octet centroid. As we have not computed the mass of the η -meson on the lattice due to the problems associated with disconnected diagrams, we have

³ The quark-mass dependence of the decuplet parameters is a higher-order effect.

used the Gell-Mann–Okubo mass relation among the square of the meson masses as input for m_η into the one-loop chiral expression in eq. (8), and used our lattice data for the π and K parameters. Expressions for the octet baryon masses at higher orders in the chiral expansion have been computed [17, 20, 21]. However, with lattice calculations at only four pion masses, and the heaviest likely outside the range of validity of the chiral theory, there is little point in attempting to fit such expressions.

As with any lattice QCD calculation, there are additional systematic uncertainties that must be estimated or removed. The finite-volume effects on the baryon masses are estimated to be negligible. The MILC lattices used in this work have spatial extent $L \sim 2.5$ fm and finite-volume chiral perturbation theory for baryons [22] indicates that for the meson masses used, the finite-volume effects are very small. The source of uncertainty in this calculation that we have the least control over is that due to the finite lattice spacing, b , however such corrections are expected to be small. The lattice spacing transforms as a singlet under flavor transformations, and consequently, it does not contribute at leading order to violations of the GMO relation. Even corrections of the form $\mathcal{O}(m_q b^2)$ will not contribute as they transform as an $\mathbf{8} \oplus \mathbf{1}$. However, we expect contributions of the form $\left(4m_K^3 - 3m_\eta^3 - m_\pi^3 \right) b^2$, which is an element of the $\mathbf{27}$. Further, there will also be exponentially-suppressed $\mathcal{O}(b)$ contributions multiplying objects that transform as a $\mathbf{27}$. The centroids of the baryon octet will experience $\mathcal{O}(b^2)$ shifts, as well as the exponentially suppressed $\mathcal{O}(b)$ contributions. A full analysis of the chiral corrections at finite lattice spacing should be performed in the mixed-action theory [23] when resources to calculate at another lattice spacing materialize. While there have been other quenched QCD [24, 25, 26, 27] and full-QCD [28, 29, 30, 31] studies of octet baryon masses, to our knowledge there have been no other full-QCD studies of the complete baryon octet in the chiral regime, prior to this work.

II. CONCLUSIONS

In this paper we have explored violations of the Gell-Mann–Okubo mass relation among baryons in the lowest-lying octet using fully-dynamical lattice QCD. With domain-wall valence quarks on rooted staggered MILC configurations, we have calculated the deviations to the GMO mass relation at four different pion masses. We find that the violations are consistent with the size of the experimental value, and vanish toward the limit of exact $SU(3)$ symmetry as required. Therefore, we conclude that the smallness of the experimentally-observed violation of the GMO mass relation is not an accident. This is consistent with the phenomenological observation that higher-dimensional representations of the flavor group are suppressed compared with lower-dimensional representations, and also consistent with the additional suppression predicted in the large- N_C limit of QCD.

Acknowledgments

We thank David Kaplan for valuable conversations and Robert Edwards for help with the QDP++/Chroma programming environment [12] with which the calculations discussed here were performed. We are also indebted to the MILC and the LHP collaborations for use of their configurations and propagators, respectively. The computations for this work were performed at the High Performance Computing Center at JLab under the auspices of the U.S. SciDAC initiative. The work of MJS is supported in part by the U.S. Dept. of En-

ergy under Grant No. DE-FG03-97ER4014. The work of KO is supported in part by the U.S. Dept. of Energy under Grant No. DF-FC02-94ER40818. The work of SRB is supported in part by the National Science Foundation under grant No. PHY-0400231. The work of KO and SRB is supported in part by DOE contract DE-AC05-84ER40150, under which the Southeastern Universities Research Association (SURA) operates the Thomas Jefferson National Accelerator Facility.

-
- [1] M. Gell-Mann, Phys. Rev. **125** (1962) 1067.
 - [2] Y. Ne'eman, Nucl. Phys. **26**, 222 (1961).
 - [3] M. Ademollo and R. Gatto, Phys. Rev. Lett. **13**, 264 (1964).
 - [4] S. Okubo, Prog. Theor. Phys. **27** (1962) 949.
 - [5] R. F. Dashen, E. Jenkins and A. V. Manohar, Phys. Rev. D **51**, 3697 (1995) [arXiv:hep-ph/9411234].
 - [6] R. G. Edwards *et al.* [LHPC Collaboration], PoS **LAT2005**, 056 (2005) [arXiv:hep-lat/0509185].
 - [7] D. B. Renner *et al.* [LHP Collaboration], Nucl. Phys. Proc. Suppl. **140**, 255 (2005) [arXiv:hep-lat/0409130].
 - [8] K. Orginos, D. Toussaint and R. L. Sugar [MILC Collaboration], Phys. Rev. D **60**, 054503 (1999) [arXiv:hep-lat/9903032].
 - [9] K. Orginos and D. Toussaint [MILC collaboration], Phys. Rev. D **59**, 014501 (1999) [arXiv:hep-lat/9805009].
 - [10] C. W. Bernard *et al.*, Phys. Rev. D **64**, 054506 (2001) [arXiv:hep-lat/0104002], <http://qcd.nersc.gov/>.
 - [11] A. Hasenfratz and F. Knechtli, Phys. Rev. D **64**, 034504 (2001) [arXiv:hep-lat/0103029].
 - [12] R. G. Edwards and B. Joo [SciDAC Collaboration], Nucl. Phys. Proc. Suppl. **140** (2005) 832 [arXiv:hep-lat/0409003].
 - [13] C. McClendon, Jlab preprint, JLAB-THY-01-29 (2001).
 - [14] E. Jenkins and A. V. Manohar, Phys. Lett. B **255**, 558 (1991).
 - [15] E. Jenkins and A. V. Manohar, UCSD-PTH-91-30 *Talk presented at the Workshop on Effective Field Theories of the Standard Model, Dobogoko, Hungary, Aug 1991*
 - [16] J. Bijnens, H. Sonoda and M. B. Wise, Nucl. Phys. B **261**, 185 (1985).
 - [17] E. Jenkins, Nucl. Phys. B **368**, 190 (1992).
 - [18] V. Bernard, N. Kaiser and U. G. Meissner, Z. Phys. C **60**, 111 (1993) [arXiv:hep-ph/9303311].
 - [19] M. N. Butler, M. J. Savage and R. P. Springer, Nucl. Phys. B **399**, 69 (1993) [arXiv:hep-ph/9211247].
 - [20] A. Walker-Loud, Nucl. Phys. A **747**, 476 (2005) [arXiv:hep-lat/0405007].
 - [21] M. Frink and U. G. Meissner, JHEP **0407**, 028 (2004) [arXiv:hep-lat/0404018].
 - [22] S. R. Beane, Phys. Rev. D **70**, 034507 (2004) [arXiv:hep-lat/0403015].
 - [23] B. C. Tiburzi, Phys. Rev. D **72**, 094501 (2005) [arXiv:hep-lat/0508019].
 - [24] T. Bhattacharya, R. Gupta and S. R. Sharpe, Nucl. Phys. Proc. Suppl. **47**, 549 (1996) [arXiv:hep-lat/9512005].
 - [25] T. Bhattacharya, R. Gupta, G. Kilcup and S. R. Sharpe, Phys. Rev. D **53**, 6486 (1996) [arXiv:hep-lat/9512021].
 - [26] K. C. Bowler *et al.* [UKQCD Collaboration], Phys. Rev. D **62**, 054506 (2000)

- [arXiv:hep-lat/9910022].
- [27] S. Aoki *et al.* [CP-PACS Collaboration], Phys. Rev. D **67**, 034503 (2003) [arXiv:hep-lat/0206009].
 - [28] C. R. Allton *et al.* [UKQCD Collaboration], Phys. Rev. D **60**, 034507 (1999) [arXiv:hep-lat/9808016].
 - [29] S. Aoki *et al.* [JLQCD Collaboration], Phys. Rev. D **68**, 054502 (2003) [arXiv:hep-lat/0212039].
 - [30] C. Aubin *et al.*, Phys. Rev. D **70**, 094505 (2004) [arXiv:hep-lat/0402030].
 - [31] Y. Namekawa *et al.* [CP-PACS Collaboration], Phys. Rev. D **70**, 074503 (2004) [arXiv:hep-lat/0404014].